

2019 ACE Value Based Measures

Two measurement periods each including one year of rolling data (with payments in December and June of each year).

2019 Pediatric Preventive

- 1- The percentage of children 2 years of age who had four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV); one measles, mumps and rubella (MMR); three haemophilus influenza type B (HiB); three hepatitis B (HepB), one chicken pox (VZV); four pneumococcal conjugate (PCV); one hepatitis A (HepA); two or three rotavirus (RV); and two influenza (flu) vaccines by their second birthday. **(NCQA)**
- 2- Patients, 12 to 21 years of age, who had at least one comprehensive well-care visit with a PCP or OB/GYN practitioner during the past year. **(NCQA)**
- 3- Children, 3–18 years of age, who were diagnosed with pharyngitis, dispensed an antibiotic and received a group A streptococcus (strep) test in the seven day period from three days prior through three days after the diagnosis of pharyngitis. **(NCQA)**
- 4- Patients, ≥ 12 years old, screened for depression in the last year with appropriate follow-up if screening is positive. **(NCQA)**
- 5- Children and adolescents 3–17 years old who had an outpatient visit with a primary care practitioner or OB/GYN during the measurement year and had evidence of body mass index (BMI) percentile documentation. **(NCQA)**
- 6- Children and adolescents 3–17 years old who had an outpatient visit with a primary care practitioner or OB/GYN during the measurement year and had evidence of counseling for nutrition. **(NCQA)**
- 7- Children and adolescents 3–17 years old who had an outpatient visit with a primary care practitioner or OB/GYN during the measurement year and had evidence of counseling for physical activity. **(NCQA)**

2019 Adult Preventive

- 1- Percentage of Medicare patients (age ≥ 20) with an annual wellness visit in the past rolling 12 months. **(NCQA)**
- 2- Patients, ≥ 12 years old, screened for depression in the last year with appropriate follow-up if screening is positive. **(NCQA)**
- 3- Patients 18 to 64 years old with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription on or three days after the episode. **(NCQA)**
- 4- **Adult patients 50–75 who had appropriate screening for colorectal cancer** with any of the following tests: annual fecal occult blood test, flexible sigmoidoscopy every 5 years, colonoscopy every 10 years, computed tomography colonography every 5 years, stool DNA test every 3 years. **(NCQA)**
- 5- Female patients, 50 through 74 years old, who had a mammogram in the last 27 months. **(NCQA)**
- 6- Patients, ≥ 18 years old, seen by the provider with a documented BMI within the last 12 months and if the BMI is outside of the normal parameters (≥ 18.5 or < 25), a follow-up is documented in the last 12 months. **(CMS)**

2019 At Risk - Diabetes

- 1- Percentage of adult patients 18-75 years of age with diabetes who had a retinal or dilated eye examination by an eye care professional in the past year, unless eye exam is normal, then an eye exam is needed every two years. **(NCQA)**
- 2- Percentage of patients 18-75 years old with diabetes who had a nephropathy screening test or evidence of nephropathy during the measurement period. **(NCQA)**
- 3- Adults 18–75 years of age with diabetes who had an HbA1c < 8 . **(NCQA)**
- 4- Adults 40-75 years of age who have diabetes and who do not have clinical ASCVD, who received statin therapy. **(NCQA)**

2019 At Risk - Cardiology

- 1- Patients with hypertension, ≥ 18 -85 years old, who had their blood pressure documented in the past year as $< 140/90$ mm Hg. **(NCQA)**
 - 2- Percentage of males 21–75 years of age and females 40–75 years of age who have clinical atherosclerotic cardiovascular disease (ASCVD) and who received statin therapy. **(NCQA)**
-